Brain Research, 567 (1991) 358–359
Elsevier

Author Index

Akita, K., see Kudo, Y., 342
Albrecht, J., see Hilgier, W., 1
Alexander, S.A., see Stea, A., 83
Alonso, M.J., Arribas, S., Marín, J., Balfagón, G. and Salaices, M., Presynaptic M_2-muscarinic receptors on noradrenergic nerve endings and endothelium-derived M_3 receptors in cat cerebral arteries, 76
Amalric, M., see Schmied, A., 91
Apicella, P., see Ljungberg, T., 337
Armstrong, D.L., see Denny, J.B., 321
Arribas, S., see Alonso, M.J., 76

Baldwin, C., see Sasek, C.A., 159
Balfagón, G., see Alonso, M.J., 76
Barbosa, J.A., see Oonnor, D.T., 188
Barkai, E., see Golan, H., 149
Baumann, M.H. and Rabii, J., Inhibition of suckling-induced prolactin release by μ- and κ-opioid antagonists, 224
Becker, D.P., see Hovda, D.A., 1
Becker, D.P., see Katayama, Y., 57
Berman, R.F., see Janusz, C.A., 181
Bieger, D., see Wang, Y.T., 260
Bonvento, G., Lacombe, P., MacKenzie, E.T. and Seylaz, J., Effects of dorsal raphe stimulation on cerebral glucose utilization in the anaesthetized rat, 325
Booze, R.M., see D'Costa, A., 204
Borde, M., Pereda, A.E. and Morales, F.R., Electrophysiological characteristics of the Mauthner cell of the weakly electric fish *Gymnotus carapo*, 145
Boyd, R.L., see D'Costa, A., 204
Bramham, C.R., Milgram, N.W. and Srebro, B., δ Opioid receptor activation is required to induce LTP of synaptic transmission in the lateral perforant path in vivo, 42
Brazitikos, P.D. and Tsacopoulos, M., Metabolic signaling between photoreceptors and glial cells in the retina of the drone (*Apis mellifera*), 33
Breese, C.R., see D'Costa, A., 204
Breese, G.R., see McCown, T.J., 25
Bruner, J., see Roux-Bruxelle, M., 175
Bruns, R.F., see Janusz, C.A., 181

Cannon, J.T., see Levine, R., 140
Caramia, M.D., see Rossini, P.M., 111
Castillo, P., Pedroarena, C., Chase, M.H. and Morales, F.R., Strychnine blockade of the non-reciprocal inhibition of trigeminal motoneurons induced by stimulation of the parvocellular reticular formation, 346
Chapin, J.K., see Nicolelis, M.A.L., 133
Chase, M.H., see Castillo, P., 346
Czternasty, J., see Roux-Bruxelle, M., 175

De Ceballos, M.L., see Pérez-Otaño, I., 127
Deminière, J.M., see Piazza, P.V., 169
Denny, J.B., Polan-Curtain, J., Wayner, M.J. and Armstrong, D.L., Angiotensin II blocks hippocampal long-term potentiation, 321
Desiato, M.T., see Rossini, P.M., 111
Dewar, D., see Ikeda, M., 51
Dormont, J.F., see Schmied, A., 91
Duncan, G.E., see McCown, T.J., 25
Durand, D., see Nakagawa, M., 241
D'Costa, A., Breese, C.R., Boyd, R.L., Booze, R.M. and Sonntag, W.E., Attenuation of Fos-like immunoreactivity induced by a single electroconvulsive shock in brains of aging mice, 204

Ehret, G. and Fischer, R., Neuronal activity and tonotopy in the auditory system visualized by *c-fos* gene expression, 350
Erickson, J.T. and Millhorn, D.E., Fos-like protein is induced in neurons of the medulla oblongata after stimulation of the carotid sinus nerve in awake and anesthetized rats, 11

Farin, D., see Schmied, A., 91
Finch, C.E., see Pasinetti, G.M., 253
Finch, C.E., see Uenishi, N., 248
Fischer, R., see Ehret, G., 350
Fujiwara, K., see Sato, K., 64

Gelhard, R.E., see Johnson, A.E., 332
Golan, H., Barkai, E. and Grossman, Y., High CO_2-bicarbonate buffer modifies GABAergic inhibitory effect at the crayfish neuromuscular synapse, 149
Grossman, Y., see Golan, H., 149
Guilbaud, G., see Kayser, V., 197
Gullans, S.R., see Verbalis, J.G., 274

Hamdi, A. and Prasad, C., Attenuation of pulsatile changes in the density of striatal [^{3}H]GBR-12935 binding sites during chronic ethanol consumption, 71
Harbaugh, C.R., see Johnson, A.E., 332
Haugvicova, R., see Hilgier, W., 1
Hazrati, L.-N. and Parent, A., Contralateral pallidothalamic and pallidotegmental projections in primates: an anterograde and retrograde labeling study, 212
Herrero, M.T., see Pérez-Otaño, I., 127
Hilgier, W., Haugvicova, R. and Albrecht, J., Decreased potassium-stimulated release of [^{3}H] D-aspartate from hippocampal slices distinguishes encephalopathy related to acute liver failure from that induced by simple hyperammonemia, 1
Hovda, D.A., Yoshino, A., Kawamata, T., Katayama, Y. and Becker, D.P., Diffuse prolonged depression of cerebral oxidative metabolism following concussive brain injury in the rat: a cytochrome oxidase histochemistry study, 1

Igisu, H., see Matsuoka, M., 328
Iijima, N., see Mori, T., 355
Ikeda, M., Dewar, D. and McCulloch, J., Selective reduction of [^{125}I]apamin binding sites in Alzheimer hippocampus: a quantitative autoradiographic study, 51
Inoue, N., see Matsuoka, M., 328
Ishida, M., see Kudo, Y., 342
Ishihara, K., see Sakai, N., 267

Janusz, C.A., Bruns, R.F. and Berman, R.F., Functional activity of the adenosine binding enhancer, PD 81,723, in the in vitro hippocampal slice, 181
Joh, T.H., see Wessel, T.C., 231
Johnson, A.E., Harbaugh, C.R. and Gelhard, R.E., Projections from the ventromedial nucleus of the hypothalamus contain oxytocin binding sites, 332

Kabiersch, A., see Lyrer, P., 317
Kaeser, H., see Lyrer, P., 317
Katayama, Y., see Hovda, D.A., 1
Katayama, Y., Tamura, T., Becker, D.P. and Tsubokawa, T., Calcium-dependent component of massive increase in extracellular potassium during cerebral ischemia as demonstrated by microdialysis in vivo, 57
Kawamata, T., see Hovda, D.A., 1
Kayser, V. and Guilbaud, G., Physiological relevance and time course of a tonic endogenous opioid modulation of nociceptive messages, based on the effects of naloxone in a rat model of localized hyperalgesic inflammation, 197
Kharoubi, M., see Piazza, P.V., 169
Kiss, J., see Sandor, N.T., 313
Kitabatake, K., see Mori, T., 355
Klein, R.L., see Oonnor, D.T., 188
Kohama, S.#, J. F. Reinhard Jr., #Cheng, H.W., see Pasinetti, G.M., 253
Kohriyama, K., see Matsuoka, M., 328
Kohsaka, S., see Mori, T., 355
Komure, O., see Sakai, N., 267
Kream, R.M., see Marchand, J.E., 290
Kreutzberg, G.W., see Nacimiento, W., 283
Kudo, Y., Akita, K., Ishida, M. and Shinozaki, H., A significant increase in intracellular Ca^{2+} concentration induced by (2*S*,3*R*,4*S*)-2-(carboxycyclopropyl-)glycine, a new potent NMDA agonist, in cultured rat hippocampal neurons, 342

Lacombe, P., see Bonvento, G., 325
Lajtha, A., see Sandor, N.T., 313
Landolt, H., see Lyrer, P., 317
Langemann, H., see Lyrer, P., 317
Lavaroni, F., see Rossini, P.M., 111
Lefebvre, P.P., Van De Water, T.R., Weber, T., Rogister, B. and Moonen, G., Growth factor interactions in cultures of dissociated adult acoustic ganglia: neu-

ronotrophic effects, 306
Levine, R., Morgan, M.M., Cannon, J.T. and Liebeskind, J.C., Stimulation of the periaqueductal gray matter of the rat produces a preferential ipsilateral antinociception, 140
Liebeskind, J.C., see Levine, R., 140
Lin, R.C.S., see Nicolelis, M.A.L., 133
Ljungberg, T., Apicella, P. and Schultz, W., Responses of monkey midbrain dopamine neurons during delayed alternation performance, 337
Luquin, M.R., see Pérez-Otaño, I., 127
Lyrer, P., Landolt, H., Kabiersch, A., Langemann, H. and Kaeser, H., Levels of low molecular weight scavengers in the rat brain during focal ischemia, 317

MacKenzie, E.T., see Bonvento, G., 325
Magoski, N.S., see Walz, W., 120
Marchand, J.E., Shimonaka, H. and Kream, R.M., Biochemical characterization and anatomical distribution of a major form of unamidated precursor of substance P in rat brain, 290
Marín, J., see Alonso, M.J., 76
Matsuoka, M., Igisu, H., Kohriyama, K. and Inoue, N., Suppression of neurotoxicity of ammonia by L-carnitine, 328
McCown, T.J., Duncan, G.E. and Breese, G.R., Neuroanatomical characterization of inferior collicular seizure genesis: 2-deoxyglucose and stimulation mapping, 25
McCulloch, J., see Ikeda, M., 51
McNeill, T.H., see Pasinetti, G.M., 253
Milgram, N.W., see Bramham, C.R., 42
Millhorn, D.E., see Erickson, J.T., 11
Miyamoto, Y., see Mori, T., 355
Moal, M.L., see Piazza, P.V., 169
Moonen, G., see Lefebvre, P.P., 306
Morales, F.R., see Borde, M., 145
Morales, F.R., see Castillo, P., 346
Morgan, M.M., see Levine, R., 140
Mori, T., Miyamoto, Y., Iijima, N., Kitabatake, K. and Kohsaka, S., Dissociation of neurite-promoting activity and protease-inhibiting function of α_2-macroglobulin in culture, 355

Nacimiento, A.C., see Nacimiento, W., 283
Nacimiento, W., Schoen, S.W., Nacimiento, A.C. and Kreutzberg, G.W., Cytochemistry of 5′-nucleotidase in the superior cervical ganglion of cat and guinea pig, 283
Nakagawa, M. and Durand, D., Suppression of spontaneous epileptiform activity with applied currents, 241
Neuman, R.S., see Wang, Y.T., 260
Nichols, N.R., see Uenishi, N., 248
Nicolelis, M.A.L., Chapin, J.K. and Lin, R.C.S., Neonatal whisker removal in rats stabilizes a transient projection from the auditory thalamus to the primary somatosensory cortex, 133
Nurse, C.A., see Stea, A., 83

Obeso, J.A., see Pérez-Otaño, I., 127
Oonnor, D.T., Klein, R.L., Thureson-Klein, A.K. and Barbosa, J.A., Chromogranin A: localization and stoichiometry in large dense core catecholamine storage vesicles from sympathetic nerve, 188
Oset, C., see Pérez-Otaño, I., 127

Parent, A., see Hazrati, L.-N., 212
Pasinetti, G.M., Kohama, S., Reinhard Jr., J.F., Cheng, H.W., McNeill, T.H. and Finch, C.E., Striatal responses to decortication. I. Dopaminergic and astrocytic activities, 253
Pedroarena, C., see Castillo, P., 346
Pereda, A.E., see Borde, M., 145
Pérez-Otaño, I., Herrero, M.T., Oset, C., De Ceballos, M.L., Luquin, M.R., Obeso, J.A. and Río, J.D., Extensive loss of brain dopamine and serotonin induced by chronic administration of MPTP in the marmoset, 127
Piazza, P.V., Rougé-Pont, F., Deminière, J.M., Kharoubi, M., Moal, M.L. and Simon, H., Dopaminergic activity is reduced in the prefrontal cortex and increased in the nucleus accumbens of rats predisposed to develop amphetamine self-administration, 169
Polan-Curtain, J., see Denny, J.B., 321
Prasad, C., see Hamdi, A., 71

Rabii, J., see Baumann, M.H., 224
Río, J.D., see Pérez-Otaño, I., 127
Rogister, B., see Lefebvre, P.P., 306
Rossini, P.M., Desiato, M.T., Lavaroni, F. and Caramia, M.D., Brain excitability and electroencephalographic activation: non-invasive evaluation in healthy humans via transcranial magnetic stimulation, 111
Rougé-Pont, F., see Piazza, P.V., 169
Roux-Bruxelle, M., Czternasty, J. and Bruner, J., Long-lasting decrease in the crayfish giant motoneuron soma excitability induced by Ca^{2+} influx, 175

Sakai, N., Sasa, M., Ishihara, K., Komure, O., Tanaka, C. and Takaori, S., Effects of L-threo-DOPS, a noradrenaline precursor, on the long-term potentiation in the rat hippocampal mossy fiber-CA3 region, 267
Salaices, M., see Alonso, M.J., 76
Sanders, S.K. and Shekhar, A., Blockade of $GABA_A$ receptors in the region of the anterior basolateral amygdala of rats elicits increases in heart rate and blood pressure, 101
Sandor, N.T., Zelles, T., Kiss, J., Sershen, H., Torocsik, A., Lajtha, A. and Vizi, E.S., Effect of nicotine on dopaminergic–cholinergic interaction in the striatum, 313
Sasa, M., see Sakai, N., 267
Sasek, C.A., Baldwin, C. and Zigmond, R.E., Distribution of vasoactive intestinal peptide- and peptide histidine isoleucine amide-like immunoreactive neurons and fibers in the thoracic spinal cord of the rat, 159
Sato, K., Yoshida, S., Fujiwara, K., Tada, K. and Tohyama, M., Glycine cleavage system in astrocytes, 64
Schmied, A., Farin, D., Amalric, M. and Dormont, J.F., Changes in motor performance and rubral single unit activity in cats after microinjections of serotonin into the red nucleus area, 91
Schoen, S.W., see Nacimiento, W., 283
Schultz, W., see Ljungberg, T., 337
Sershen, H., see Sandor, N.T., 313
Seylaz, J., see Bonvento, G., 325
Shekhar, A., see Sanders, S.K., 101
Shimonaka, H., see Marchand, J.E., 290
Shinozaki, H., see Kudo, Y., 342
Shors, T.J., see Uenishi, N., 248
Simon, H., see Piazza, P.V., 169
Sonntag, W.E., see D'Costa, A., 204
Srebro, B., see Bramham, C.R., 42
Stea, A., Alexander, S.A. and Nurse, C.A., Effects of pH_i and pH_e on membrane currents recorded with the perforated-patch method from cultured chemoreceptors of the rat carotid body, 83
Strahlendorf, H.K., see Wang, Y., 153
Strahlendorf, J.C., see Wang, Y., 153

Tada, K., see Sato, K., 64
Takaori, S., see Sakai, N., 267
Tamura, T., see Katayama, Y., 57
Tanaka, C., see Sakai, N., 267
Thompson, R.F., see Uenishi, N., 248
Thureson-Klein, A.K., see Oonnor, D.T., 188
Tohyama, M., see Sato, K., 64
Torocsik, A., see Sandor, N.T., 313
Tsacopoulos, M., see Brazitikos, P.D., 33
Tsubokawa, T., see Katayama, Y., 57

Uenishi, N., Shors, T.J., Finch, C.E., Nichols, N.R. and Thompson, R.F., Increased synthesis of two polypeptides in area CA1 of the hippocampus in response to repetitive electrical stimulation, 248

Van De Water, T.R., see Lefebvre, P.P., 306
Verbalis, J.G. and Gullans, S.R., Hyponatremia causes large sustained reductions in brain content of multiple organic osmolytes in rats, 274
Vizi, E.S., see Sandor, N.T., 313
Volpe, B.T., see Wessel, T.C., 231

Walz, W. and Magoski, N.S., Short-circuiting effects of K^+ currents on electrical responses of type-1-like astrocytes from mouse cerebral cortex, 120
Wang, Y., Strahlendorf, J.C. and Strahlendorf, H.K., A transient voltage-dependent outward potassium current in mammalian cerebellar purkinje cells, 153
Wang, Y.T., Bieger, D. and Neuman, R.S., Activation of NMDA receptors is necessary for fast information transfer at brainstem vagal motoneurons, 260
Wayner, M.J., see Denny, J.B., 321
Weber, T., see Lefebvre, P.P., 306
Wessel, T.C., Joh, T.H. and Volpe, B.T., In situ hybridization analysis of c-*fos* and c-*jun* expression in the rat brain following transient forebrain ischemia, 231

Yoshida, S., see Sato, K., 64
Yoshino, A., see Hovda, D.A., 1

Zelles, T., see Sandor, N.T., 313
Zigmond, R.E., see Sasek, C.A., 159